arXiv:1203.2122v1 [math.PR] 9 Mar 2012

Stirling's approximation for central polynomial coefficients

Steffen Eger

Abstract

We derive asymptotic formulae for central polynomial coefficients, a generalization of binomial coefficients, using the distribution of the sum of independent uniform random variables and the CLT.

1 Stirling's formula and binomial coefficients

For a nonnegative integer m, Stirling's formula

$$m! \sim \sqrt{2\pi m}\Big(\frac{m}{e}\Big)^m,$$

where e is Euler's number, yields an approximation of the central binomial coefficient $\binom{m}{m/2}$ (assuming that m is even) using $\binom{m}{k} = \frac{m!}{k!(m-k)!}$ as

$$\binom{m}{m/2} \sim \frac{2^{m+1}}{\sqrt{2\pi m}},$$

where we write $a_m \sim b_m$ as a short-hand for $\lim_{m\to\infty} \frac{a_m}{b_m} = 1$. In the current note, we derive an asymptotic formula for the central *multinomial triangle* or, *polynomial*, *coefficient* (cf. [1], [2]), where multinomial triangles are a generalization of binomial triangles, where entries in row k are defined as coefficients of the polynomial $(1 + x + x^2 + \ldots + x^l)^k$ for $l \geq 0$. Our derivation is not based upon asymptotics of factorials but upon the limiting distribution of the sum of discrete uniform random variables.[1]

Polynomial coefficients are important, for example, because they denote the number of *integer compositions* of the nonnegative integer n with parts in the set $A := \{a, a+1, \ldots, b\}$, where a, b, $0 \leq a \leq b$, are nonnegative integers (cf. [3]), i.e. the number of possibilities to write n as a sum of integers $p_1, \ldots, p_k$, with $p_i \in A$, for $i = 1, \ldots, k$.

2 Multinomial triangles

In generalization to binomial triangles, $(l+1)$-nomial triangles, $l \geq 0$, are defined in the following way. Starting with a 1 in row zero, construct an entry in row k, $k \geq 1$, by adding the overlying $(l+1)$ entries in row $(k-1)$ (some of these entries are taken as zero if not defined); thereby, row k has $(kl+1)$ entries. For example, the monomial ($l = 0$), binomial ($l = 1$), trinomial ($l = 2$) and quadrinomial triangles ($l = 3$) start as follows,

$$\begin{array}{c} 1 \\ 1 \\ 1 \\ 1 \end{array} \qquad \begin{array}{cccc} 1 & & & \\ 1 & 1 & & \\ 1 & 2 & 1 & \\ 1 & 3 & 3 & 1 \end{array} \qquad \begin{array}{ccccccc} 1 & & & & & & \\ 1 & 1 & 1 & & & & \\ 1 & 2 & 3 & 2 & 1 & & \\ 1 & 3 & 6 & 7 & 6 & 3 & 1 \end{array} \qquad \begin{array}{cccccccccc} 1 & & & & & & & & & \\ 1 & 1 & 1 & 1 & & & & & & \\ 1 & 2 & 3 & 4 & 3 & 2 & 1 & & & \\ 1 & 3 & 6 & 10 & 12 & 12 & 10 & 6 & 3 & 1 \end{array}$$

---

[1] Throughout, we assume that m or l are even. If this is not the case, replace respective quantities, e.g. $\frac{ml}{2}$, with their floor, $\lfloor \frac{ml}{2} \rfloor$.

In the $(l+1)$-nomial triangle, entry n, $0 \leq n \leq kl$, in row k, which we denote by $\binom{k}{n}_{l+1}$, has the following interpretation. It is the coefficient of x^n in the expansion of

$$(1 + x + x^2 + \ldots + x^l)^k = \sum_{n=0}^{kl} \binom{k}{n}_{l+1} x^n. \tag{2.1}$$

In recent work, Eger [3] has shown that $\binom{k}{n}_{l+1}$ denotes the number of integer compositions of the nonnegative integer n with k parts $p_1, \ldots, p_k$, each from the set $\{0, 1, \ldots, l\}$, and allows the following representation,

$$\binom{k}{n}_{l+1} = \sum_{\substack{k_0+\ldots+k_l=k \\ 0\cdot k_0+1\cdot k_1+\ldots+l\cdot k_l=n}} \binom{k}{k_0, \ldots, k_l}, \tag{2.2}$$

where $\binom{k}{k_0, \ldots, k_l}$ is a *multinomial coefficient*, defined as $\frac{k!}{k_0! \ldots k_l!}$, for nonnegative integers $k_0, \ldots, k_l$. We can easily verify representation (2.2) by noting that for real numbers $x_0, \ldots, x_l$, it holds that (cf. [5])

$$(x_0 + x_1 + \ldots + x_l)^k = \sum_{\substack{k_0,\ldots,k_l \geq 0 \\ k_0+\ldots+k_l=k}} \binom{k}{k_0, \ldots, k_l} x_0^{k_0} \ldots x_l^{k_l}.$$

Then setting $x_i = x^i$ for $i = 0, \ldots, l$,

$$(1 + x + x^2 + \ldots + x^l)^k = \sum_{\substack{k_0,\ldots,k_l \geq 0 \\ k_0+\ldots+k_l=k}} \binom{k}{k_0, \ldots, k_l} x^{0\cdot k_0+\ldots+l\cdot k_l}, \tag{2.3}$$

and comparing coefficients of the right-hand sides of (2.1) and (2.3) leads to (2.2).

3 Generalized Stirling's formula

Our strategy for deriving approximation formulae for polynomial coefficients is as follows. First, we determine the asymptotic distribution of the sum of discrete uniform variables, which we easily find to be a normal distribution by the Central Limit Theorem (CLT). Then we determine the exact distribution, which turns out to yield the normalized polynomial coefficients $\binom{m}{n}_{l+1}$. This distribution has recently been found by Caiado and Rathie [1] using characteristic functions, residue theory and Cauchy's theorem, but we give an elementary two-line proof that almost requires no effort at all. By relating the asymptotic distribution to the exact distribution, we obtain a multinomial analogue of Stirling's approximation to the central binomial coefficient.

3.1 Distribution of the sum of discrete uniform variables

Let m be a positive integer and let l be a nonnegative integer. Let S_j, $j = 1, \ldots, m$, be identically and independently distributed random draws from the discrete uniform distribution on the set $\{0, \ldots, l\}$, and let $S^{(m)}$ be their sum,

$$S^{(m)} = \sum_{j=1}^{m} S_j.$$

Obviously, the mean and variance of $S^{(m)}$ are given by

$$\mu_{m,l} := \mathrm{E}[S^{(m)}] = m\,\mathrm{E}[S_j] = \frac{ml}{2}, \qquad \sigma_{m,l}^2 := \mathrm{Var}[S^{(m)}] = m\,\mathrm{Var}[S_j] = m\frac{(l+1)^2-1}{12},$$

by standard moments of the uniform distribution. Moreover, by the CLT

$$\sqrt{m}\Big(\frac{S^{(m)}}{m} - \frac{l}{2}\Big) \rightarrow \mathcal{N}(0, \frac{(l+1)^2-1}{12}) \quad \text{as } m \rightarrow \infty,$$

so that for large m, $S^{(m)}$ is approximately normally distributed with mean $\frac{ml}{2}$ and variance $m\frac{(l+1)^2-1}{12}$.

However, we can also exactly determine the probability that $S^{(m)}$ takes on the integer value n, for $0 \leq n \leq ml$. For this purpose, interpret all S_j, $j = 1, \dots, m$, as independently and identically *multinomially* distributed with probabilities $p_0 = \dots = p_l = \frac{1}{l+1}$ of 'events' 0 to l occurring, i.e. each S_j is vector-valued, $S_j = (A_0, \dots, A_l)$, with $P[S_j = (a_0, \dots, a_l)] = \frac{1}{l+1}$ for $a_0 + \dots + a_l = 1$, $a_k \geq 0$, where A_k denotes the number of times event k occurs, for $k = 0, \dots, l$. Hence, the sum $S^{(m)}$ is multinomially distributed with probability mass function

$$P[S^{(m)} = (a_0, \dots, a_l)] = P[A_0 = a_0, \dots, A_l = a_l] = \binom{m}{a_0, \dots, a_l}\Big(\frac{1}{l+1}\Big)^m,$$

where $a_0 + \dots + a_l = m$. Then, if $S^{(m)} = (a_0, \dots, a_l)$ for nonnegative integers $a_0, \dots, a_l$ with $a_0 + \dots + a_l = m$, $S^{(m)}$ 'represents' the integer $0 \cdot a_0 + \dots + l \cdot a_l$. Thus, for n such that $0 \leq n \leq ml$, we obviously have, reinterpreting $S^{(m)}$ as scalar-valued,

$$P[S^{(m)} = n] = \sum_{\substack{a_0+\dots+a_l=m \\ 0\cdot a_0+\dots+l\cdot a_l=n}} P[S^{(m)} = (a_0, \dots, a_l)] = \Big(\frac{1}{l+1}\Big)^m \sum_{\substack{a_0+\dots,a_l=m \\ 0\cdot a_0+\dots+l\cdot a_l=n}} \binom{m}{a_0, \dots, a_l} = \Big(\frac{1}{l+1}\Big)^m \binom{m}{n}_{l+1}.$$

3.2 Approximation Formulae

From our above derivations, we know that the normalized random variable $S^{(m)}$ converges in distribution to a normally distributed random variable, which precisely means that the distribution function of $S^{(m)}$ converges pointwise to the distribution function of the corresponding normally distributed random variable. Let $0 \leq n \leq ml$ and consider

$$P[S^{(m)} = n] = P[S^{(m)} \leq n] - P[S^{(m)} \leq n-1].$$

By convergence of distribution functions, we hence know that, for large m,

$$P[S^{(m)} = n] \approx P[n-1 \leq X \leq n] \approx P[n-1/2 \leq X \leq n+1/2], \tag{3.1}$$

where X has a normal distribution with the same mean and variance as $S^{(m)}$, and applying a continuity correction (cf. [4]). Rewriting the right hand side of (3.1) in terms of the standard normal variable $Z = \frac{X-\mu_{m,l}}{\sigma_{m,l}}$, we thus obtain for $n = \mu_{m,l}$,

$$P[S^{(m)} = \frac{ml}{2}] \approx P[-\frac{1}{2\sigma_{m,l}} \leq Z \leq +\frac{1}{2\sigma_{m,l}}] = \Phi\left(\frac{1}{2\sigma_{m,l}}\right) - \Phi\left(-\frac{1}{2\sigma_{m,l}}\right),$$

where Φ is the standard normal cumulative distribution function. Geometrically, $\Phi\left(\frac{1}{2\sigma_{m,l}}\right) - \Phi\left(-\frac{1}{2\sigma_{m,l}}\right)$ denotes the area under the normal curve $\phi(z) = 1/\sqrt{2\pi}\exp(-z^2/2)$ from $-\frac{1}{2\sigma_{m,l}}$ to $+\frac{1}{2\sigma_{m,l}}$. For large m, this area can be approximated by the area of the rectangle with width $\frac{1}{\sigma_{m,l}}$ and height $\phi(0)$, that is,

$$P[S^{(m)} = \frac{ml}{2}] \approx \frac{1}{\sqrt{2\pi m\frac{(l+1)^2-1}{12}}}.$$

Using the expression for $P[S^{(m)}=n]$ from above, we hence have

$$\binom{m}{\frac{ml}{2}}_{l+1} \sim \frac{(l+1)^m}{\sqrt{2\pi m \frac{(l+1)^2-1}{12}}}.$$

For $l=1$, Pascal's case, $l=2$, $l=3$, and $l=4$, we therefore have the approximations

$$\binom{m}{\frac{m}{2}} \sim \frac{2^{m+1}}{\sqrt{2\pi m}}, \qquad \binom{m}{m}_3 \sim \frac{3^m}{\sqrt{\frac{4}{3}\pi m}}, \qquad \binom{m}{\frac{3}{2}m}_4 \sim \frac{4^m}{\sqrt{\frac{5}{2}\pi m}}, \qquad \binom{m}{2m}_5 \sim \frac{5^m}{2\sqrt{\pi m}}.$$



Figure 1: Distributions $P[S^{(m)}=n]$ for $m=5,10,20$ for $l=4$ fixed; and normal approximations.

In Figure 3.2, we show for $l=4$ the distributions $P[S^{(m)}=n]$ for $m=5,10,20$ and their respective normal approximations.

These plots suggest to more generally consider a pointwise approximation of $P[S^{(m)}=n]$, for $0\le n\le ml$, by the value $f(n)$ of the density of the corresponding normally distributed random variable X, leading to the following approximation formulae for polynomial coefficients $\binom{m}{n}_{l+1}$,

$$\binom{m}{n}_{l+1} \approx \frac{(l+1)^m}{\sqrt{2\pi\sigma_{m,l}^2}}\exp\left(-\frac{(n-\mu_{m,l})^2}{2\sigma_{m,l}^2}\right).$$

How well are these 'normal approximations' $\hat{a}_{m,n,l}$ of $a_{m,n,l}:=\binom{m}{n}_{l+1}$ for various values of n for finite m? In Figure 3.2, we depict the relative error $\left|\frac{\hat{a}_{m,n,l}-a_{m,n,l}}{a_{m,n,l}}\right|$ for all values of n between 0 and ml for $m=100,200$ and $l=3,4$. We see that the relative error becomes very low for n 'around' $\frac{ml}{2}$ and increases symmetrically as n approaches the boundary values 0 and ml. In the same figure, we show the decrease of the relative error for the central polynomial coefficient as a function of m.

References

[1] Caiado, C.C.S., and Rathie, P.N. (2007). Polynomial Coefficients and Distribution of the Sum of Discrete Uniform Variables, in Mathai, A. M., Pathan, M. A. Jose, K. K. and Jacob, J., eds., *Eighth Annual Conference of the Society of Special Functions and their Applications*, Pala, India, Society for Special Functions and their Applications.

[2] Comtet, L. (1974). *Advanced Combinatorics*, D. Reidel Publishing Company.

[3] Eger, S. (2011). Integer compositions and Multinomial Triangles, submitted.

[4] Walsh, D.P. (1995). Equating Poisson and Normal Probability Functions to derive Stirling's formula. *The American Statistician*, 49(3): 270–271.

[5] Weisstein, Eric W. "Multinomial Series." From MathWorld – A Wolfram Web Resource. http://mathworld.wolfram.com/MultinomialSeries.html



Figure 2: Left: Relative error of normal approximation of $\binom{m}{n}_{l+1}$ for $m = 100, 200$ and $l = 3, 4$ as a function of n. Right: Relative error for central polynomial coefficient as a function of m for $l = 4$.